

Ocean warming and acidification have complex interactive effects on the dynamics of a marine fungal disease

Williams, G.J.; Price, N.N.; Ushijima, B.; Aeby, G.S.; Callahan, S.; Davy, S.K.; Gove, J.M.; Johnson, M.D.; Knapp, I.S.; Shore-Maggio, A.; Smith, J.E.; Videau, P.; Work, T.M.

Proceedings of the Royal Society B: Biological Sciences

DOI:

[10.1098/rspb.2013.3069](https://doi.org/10.1098/rspb.2013.3069)

Published: 07/03/2014

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Williams, G. J., Price, N. N., Ushijima, B., Aeby, G. S., Callahan, S., Davy, S. K., Gove, J. M., Johnson, M. D., Knapp, I. S., Shore-Maggio, A., Smith, J. E., Videau, P., & Work, T. M. (2014). Ocean warming and acidification have complex interactive effects on the dynamics of a marine fungal disease. *Proceedings of the Royal Society B: Biological Sciences*, 281(1778). <https://doi.org/10.1098/rspb.2013.3069>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Ocean warming and acidification have complex interactive effects on the dynamics of a marine fungal disease

Running title: CCA fungal disease dynamics

Gareth J. Williams^{1*§}, Nichole N. Price^{1§}, Blake Ushijima^{2,3}, Greta S. Aeby³, Sean Callahan², Simon K. Davy⁴, Jamison M. Gove^{5,6}, Maggie D. Johnson¹, Ingrid S. Knapp⁴, Amanda Shore-Maggio^{2,3}, Jennifer E. Smith¹, Patrick Videau², Thierry M. Work⁷

¹Scripps Institution of Oceanography, Center for Marine Biodiversity and Conservation, University of California San Diego, La Jolla, California, 92093, USA.

²University of Hawai'i at Mānoa, Department of Microbiology, Honolulu, HI, 96822, USA

³Hawai'i Institute of Marine Biology, Kaneohe, HI 96744, USA.

⁴School of Biological Sciences, Victoria University of Wellington, PO Box 600, Wellington, New Zealand.

⁵Coral Reef Ecosystem Division (CRED), Pacific Islands Fisheries Science Center (PIFSC), NOAA, 1610 Kapiolani Blvd., Suite 1110, Honolulu, HI 96814, USA.

⁶Joint Institute for Marine and Atmospheric Research, University of Hawaii at Manoa, Honolulu, Hawaii, USA

⁷US Geological Survey, National Wildlife Health Center, Honolulu Field Station, PO Box 50167, Honolulu, HI 96850, USA.

*corresponding author, email: gareth@ucsd.edu, Tel: 858-534-8783, Fax: 858-822-1267

§authors contributed equally to this work

Abstract: Diseases threaten the structure and function of marine ecosystems and are contributing to the global decline of coral reefs. We currently lack an understanding of how climate change stressors, such as ocean acidification and warming, may simultaneously affect coral reef disease dynamics, particularly diseases threatening key reef-building organisms such as crustose coralline algae (CCA). Here we use coralline fungal disease (CFD), a previously described CCA disease from the Pacific, to examine these simultaneous effects using both field observations and experimental manipulations. We identify the associated fungus as belonging to the subphylum *Ustilaginomycetes* and show linear lesion expansion rates on individual hosts can reach 6.5mm per day. Further, we demonstrate for the first time that ocean warming events could increase the frequency of CFD outbreaks on coral reefs, but that ocean acidification-induced lowering of pH may ameliorate outbreaks by slowing lesion expansion rates on individual hosts. Lowered pH may still reduce overall host survivorship, however, by reducing calcification and facilitating fungal bioerosion. Such complex, interactive effects between simultaneous extrinsic environmental stressors on disease dynamics are important to consider if we are to accurately predict the response of coral reef communities to future climate change.

Keywords: coral reef; coralline fungal disease; ocean acidification; temperature; bioerosion; climate change

1. INTRODUCTION

Diseases alter ecosystems [1] and threaten marine community function and resilience [2]. On coral reefs, disease outbreaks are considered a key contributor to the recent global decline of reef health and resilience[3]. Both global impacts, such as sea-surface temperature anomalies, and local human impacts, such as pollution, drive disease dynamics and outbreaks in scleractinian corals on reefs [4, 5]. These stressors likely increase pathogen virulence and reduce host resistance, enhancing disease establishment and progression [2, 6]. Our understanding of diseases that threaten other key calcifying (reef-building) organisms, however, is rudimentary. Crustose coralline algae (CCA) serve essential functional roles in coral reef ecosystems, including facilitating reef accretion and consolidation [7], providing a settlement substrate for coral larvae [8], and forming a key successional state promoting reef recovery following acute disturbance [9]. While CCA can occupy up to 50 % of the living reef benthos [7, 10], relatively little is known about their biology and ecology [11], particularly their susceptibility to and subsequent impacts from disease [12].

Diseases can cause drastic reductions in CCA populations on coral reefs, with knock-on effects that promote regime shifts to fleshy macroalgal dominance and loss of functional resilience [13-15]. Several CCA diseases have been documented, although almost nothing is known about their etiology, spatiotemporal dynamics, and relationships with extrinsic environmental drivers [12]. This information is essential if we are to actively manage CCA disease occurrence and mitigate outbreaks at a local scale on coral reefs. While CCA are able to photoacclimate [16], they are still vulnerable to bleaching as a result of increased temperature, and their calcification, photophysiology, and survival are threatened by a lowering of pH and carbonate saturation state (Ω) as a result of ocean acidification (OA) [17-19]. However, the influence of these global-scale

stressors on local-scale CCA disease dynamics and occurrence on coral reefs is virtually unknown. Here, using a CCA disease previously described from the South Pacific, we shed light on this urgent research priority for the first time.

Coralline fungal disease (CFD) was first observed in 1997 on shallow (< 20 m) reef habitats in American Samoa. Based on gross morphology, the etiology of the disease was identified as an undescribed fungal pathogen [20]. Since 1997, CFD has been documented throughout other parts of the Pacific, with high prevalence at remote islands in the Central Pacific, in particular at Kingman Reef and Palmyra Atoll in the northern Line Islands [12]. While the distribution of CFD throughout the Pacific appears to be highly variable, evidence suggests that variations in sea-surface temperature (SST) may, in part, be driving spatial variation in disease occurrence, with higher mean SST at islands correlating with higher CFD occurrence [12]. However, the principal environmental drivers of temporal variation in CFD occurrence remain unknown.

In late 2009, a well-developed El Niño resulted in anomalous ocean warming across the equatorial Pacific Ocean. The sustained increase in temperature resulted in coral bleaching events and disease outbreaks at some of the Pacific's most remote coral reef systems, including Palmyra Atoll, where CFD is prevalent [21-23]. Using time-series field observations at Palmyra before, during, and after the 2009-2010 El Niño ocean warming event, followed by experimental manipulations, we show that CFD temporal dynamics are driven by variations in seawater temperature and that spatial variation in disease occurrence appears independent of host abundance. We show that CFD can exhibit very rapid progression across hosts, rivalling known rates for coral diseases, and that increased temperature accelerates CFD lesion progression. We show that these effects are counteracted by a reduction in pH; however, under simulated ocean acidification conditions, diseased hosts still experience greater rates of net dissolution than

healthy individuals. These complex interactions highlight the challenges associated with predicting disease outbreaks and their dynamics in a changing climate.

2. MATERIALS AND METHODS

(a) *Histopathology*

Fragments ($n = 7$, $\sim 3 \text{ cm}^2$) of crustose coralline algae (CCA) displaying gross signs of coralline fungal disease (CFD) were collected at Palmyra Atoll ($05^\circ 52' \text{ N}$, $162^\circ 06' \text{ W}$), at 10 m depth on the forereef (Fig. 1). Each sample was fixed in zinc-formaldehyde solution (Z-fix, Anatech) diluted 1:5 with ambient non-filtered seawater, decalcified using a formic acid/formaldehyde solution (Cal-Ex II, Fisher Scientific), embedded in paraffin, sectioned at $5 \mu\text{m}$, and stained with hematoxylin and eosin. Grocott's methenamine silver was used to confirm presence of fungal hyphae.

(b) *Culture conditions and phylogenetic determination of CFD fungus*

Culture conditions: To culture the fungus associated with CFD, Medium B, Wickerham's, Czapek-Dox agar, and cellulose agar [24] prepared with 25, 50, 75 or 100 % FSW as a base was used. Media were prepared with and without inclusion of the antibiotics ampicillin and spectinomycin at a concentration of $100 \mu\text{g/ml}$. Media with antibiotics was used to suppress bacterial growth, and the same media were used without antibiotics to preclude any potential negative effects on fungal growth. Solid media included 0.7, 1.0, 1.2, or 1.5 % agar (Fisher Scientific). Cultures were incubated at 25, 27, or 30°C with aeration (solid cultures) or without aeration (liquid cultures) for 21 days.

DNA extraction and gene amplification: Frozen, ground CFD lesions were thawed on ice, $5 \mu\text{l}$ were spread on PALM PEN membrane slides (Carl Zeiss), and individual fungal filaments were isolated with a Zeiss PALM laser microdissection system. DNA was extracted from filaments as

in [25] with the following modifications: Two samples of 8-10 fungal filaments in 50 µl of sterile water were lysed with a mini-beadbeater (Biospec) and extracted with phenol-chloroform. DNA was precipitated, dried, and re-suspended in 20 µl of sterile 10 mM Tris buffer, pH 8.5. A 719 bp fragment of the 18S rRNA gene was PCR amplified with fungus-specific primers designed from mixed environmental samples: nu-SSU-0817-5' and nu-SSU-1536-3' [26] and Phusion High-fidelity Polymerase (New England Biolabs). The same primers were used for sequencing.

Phylogenetic analysis: The fungal 18S rRNA gene sequence was initially assessed using NCBI and the BLAST algorithm. To further assess its relatedness to other phylum *Basidiomycota* members, especially the subphylum *Ustilaginomycetes*, 653 bp 18S rRNA gene sequences were aligned in MEGA5 [27]. The fungal isolate's sequence was aligned with 22 other NCBI sequences, chosen based on a previously published analysis of the phylum *Basidiomycota* [28], using ClustalW. A phylogenetic tree was constructed using the neighbour-joining method with 1000 bootstrap replicates [29]. All positions containing gaps and missing data were eliminated, leaving a total of 647 positions. Evolutionary distances (number of base substitutions per site) were computed using the Maximum Composite Likelihood method. Rate variation among sites was modelled with a gamma distribution (shape parameter = 8). Evolutionary analyses were conducted in MEGA5.

(c) *CFD occurrence, host cover, and associated changes in seawater temperature*

To quantify CFD occurrence, 59 × 200 m² transects were surveyed in Jul-Aug and Oct-Nov 2008 (total of 11,800 m² of reef). Backreef (n = 24 transects, 1-5 m depth), reef terrace (n = 5, 4-5 m depth), and forereef (n = 30, 10 m depth) habitats were surveyed within 12 permanent sites (Fig.

1). Along each transect, percent CCA cover was estimated using the photoquadrat method. Each photoquadrat ($n = 20$ per 50 m) was 0.63 m^2 . Percent cover was calculated *post-hoc* by identifying 100 points in a stratified random design for each photograph and averaging within each transect. Difficulties with delineating individual CCA crusts *in situ* meant a true CFD prevalence (proportion of individuals displaying signs of the disease) could not be calculated. Instead, the numbers of CFD cases were normalized to host cover (per m^2 of CCA) along each transect (see Supplemental Table 1 for mean raw number of CFD cases and percent cover of CCA within each transect over time). Forty of the 59 transects surveyed in 2008 became permanent as part of the Palmyra disease monitoring program and were re-surveyed for CFD occurrence and CCA cover in Oct-Nov 2009 and Mar 2010.

Throughout the study, *in situ* forereef temperature data were sampled at 30 min intervals at 10 m depth using Sea-Bird Electronics (SBE) 39 temperature sensors with a resolution and accuracy of 0.0001°C and $\pm 0.002^\circ\text{C}$, respectively. Satellite-derived weekly sea surface temperature (SST) and long-term monthly climatological SST for Palmyra were calculated following [30].

(d) CFD vital rates during the El Niño

On the central south forereef, 13 CFD cases were photographed weekly for four weeks in Oct-Nov 2009. Individual CFD cases were initially marked with a stainless steel pin as a reference point. Photographs were taken perpendicular to the substrate to minimize angle variations among images. CFD vital rates (lesion surface area and linear progression rate) were calculated *post-hoc* using ImageJ (<http://rsbweb.nih.gov/ij>). Active lesion surface area was the area displaying fungal cover (blue-black discolouration), not including dead CCA left in the lesion's path. Calculations

were averaged across three images of each lesion to account for slight variations in angle among photos.

(e) *Temperature and acidification experiments*

We used a factorial CO₂ bubbling and heating experiment (June 3rd to 13th 2012) to examine independent and interactive effects of ocean acidification (OA) and warming on CFD disease dynamics (lesion surface area and linear progression rate) and CCA growth (net calcification). Samples were collected, as for histopathology, from independent diseased and healthy CCA crusts. Epiphytes were removed, and each CCA genus was confirmed using a dissecting microscope. Paired CCA fragments (~2 cm² *Neogoniolithon* sp.; diseased and healthy) were placed in 1L glass aquaria holding fresh seawater equilibrated to treatment conditions (~24 – 48 h before) [sensu 31]. Experimental OA conditions were created by bubbling pre-mixed air enriched with excess *p*CO₂ (AirGas Pro) to 1124 ± 88 μ atm (mean \pm SE hereafter) to reflect atmospheric CO₂ concentrations projected in 2100 [scenario IV; 32]. In control aquaria, present-day CO₂ conditions were created using a Pacbrake 12V HP625 Air Compressor delivering ambient air. Aquaria were immersed in flow-through water baths at 28.06 ± 0.01 °C (seasonal average for Palmyra; ambient conditions) or 29.49 ± 0.02 °C (mean SST during the El Niño warming event at Palmyra; warming conditions). Four independent water baths (two ambient and two warmed) held experimental aquaria that were randomly assigned to elevated or present-day CO₂ conditions to create every combination of warming and OA treatment level (*n* = 12 per level divided evenly among the two water baths).

Aquaria were covered to prevent evaporation and rainwater from affecting salinity and placed under a shade cloth to mimic the natural light environment at 10 m on the forereef. To control for

algal metabolism, two empty aquaria per water bath were subjected to the same four treatments described above. In each aquarium, temperature and light conditions were recorded every 15 min using Onset HOBO® Pendant UA-002-64 light and temperature loggers (Suppl. Fig. 1). Light intensities (lux) were converted to availability of photosynthetically active radiation (PAR) using the equation: $1 \mu\text{mol quanta (400 – 700 nm) m/s} = 51.2 \text{ lux}$ [sensu 33]. These conversions were validated by midday PAR measurements with a Li-Cor LI 192 4π quantum sensor ($492 \pm 25 \mu\text{mol photons m/s}$). Once daily, $\text{pH}_{\text{seawater}}$ (resolution ± 0.01), temperature ($\pm 0.1 \text{ }^{\circ}\text{C}$), and dissolved oxygen ($\pm 0.2 \text{ mg/L}$, Suppl. Table 2) were monitored using a HACH HQ 40d handheld meter. These measurements were also taken at 0600h, 1200h, 1800h, and 2400h to quantify diurnal fluctuations in each aquarium (Suppl. Table 2). Twelve water samples for total alkalinity (A_T), total dissolved inorganic carbon (C_T), and salinity were collected at days 2, 6, and 10 of the experiment in 500 mL Corning brand Pyrex bottles and fixed with 200 μL saturated HgCl_2 solution (1 % headspace). Samples were collected (in duplicate) from experimental aquaria and control (empty) aquaria at each of the four treatment levels.

(f) CCA calcification and disease progression

Net calcification rates of all fragments of *Neogoniolithon* sp. were quantified with the buoyant weight method [34] (to the nearest mg) using a weigh-below basket on a balance (Denver SI-403). Changes in buoyant weight over the 10-day experiment, normalized to initial fragment mass, approximated net calcification rate. For each diseased fragment, CFD disease vital rates (lesion surface area and linear progression rate) were calculated using ImageJ as described above.

(g) Experimental water chemistry analysis

Carbonate chemistry and salinity were analysed in the Dickson Lab at Scripps Institution of Oceanography. C_T was determined using a Single Operator Multi-parameter Metabolic Analyzer (SOMMA) and a UIC Model 5011 CO_2 coulometer. A_T was determined by open-cell acid titration using a Metrohm Dosimat Model 665 and Metrohm potentiometric pH probe and meter. Salinity was determined using a Mettler Toledo Model DE45 density meter. Seawater dissolved inorganic carbon parameters (HCO_3^- , CO_3^{2-} , CO_2 , pCO_2) and saturation state of carbonate minerals (Ω -calcite and Ω -Mg calcite) were calculated based on measured C_T and A_T using the computer program Seacarb [35] and stoichiometric dissociation constants [36] (Suppl. Table 3).

(h) Data analyses

To test for differences in CCA cover and CFD occurrence across forereef sites in 2008, we ran a permutation-based analysis of variance and subsequent pairwise comparisons using PERMANOVA+ [37]. To test for any relation between CCA abundance and CFD occurrence in 2008, we used a permutational linear model using *Distlm_forward* [38]. Two-way nested analyses of variance (ANOVAs) tested whether OA (fixed) and warming (fixed) treatments independently or jointly affected net calcification; each fixed factor had two levels. Replicates from a water bath were nested within temperature treatments to test for location bias. Analyses were run independently for diseased and healthy samples. Normality and homoscedasticity were verified using the Shapiro-Wilk test. Growth responses were compared between diseased and healthy CCA within each treatment using t-tests. Proportional changes in lesion area were analysed using the Dunn's method for joint ranking, a non-parametric approach that compares means of treatments against a control (ambient SST and air); we confirmed that variances across treatments were equal with a Brown-Forsythe test. Unless otherwise stated, all analyses were completed using R 2.15.2 (R Development Core Team, <http://www.r-project.org>).

3. RESULTS

(a) *CFD gross morphology and histopathology; phylogeny of associated fungus*

CFD lesions were characterized by a diffuse area of mottled white discolouration separated from a pink CCA thallus by a blue-black band (approximately 1-3 cm wide) with irregular distinct undulating borders (Fig. 2a). Septated hyphae with branches originating near septal junctions were commonly observed (Fig. 2b). Many of the fungal filaments had highly branched structures resembling conidiophores, which in some cases appeared to have attached spherical conidia, the asexual spores of fungi. These were the dominant fungal structures in diseased CCA but were absent in healthy CCA. The CCA cuticle was overlaid and disrupted by mats of brown to colourless filamentous branching septate structures (Fig. 2c). These irregularly walled structures infiltrated into the CCA thallus at right angles to the cuticle to a depth of ~100 µm. These were morphologically compatible with fungi, and Grocott's methenamine silver confirmed a fungal infection (Fig. 2c).

Isolation by microdissection followed by genetic sequencing of the fungal hyphae revealed that 653 bp of the 18S rRNA gene sequence (Accession #KF255580) shared 97-98 % sequence identity with members of the phylum *Basidiomycota* and uncultured marine isolates. The sequence was most similar (98 % sequence identity) to members of the subphylum *Ustilaginomycetes*, in particular *Malassezia restricta* strain CBS-7877 and the marine isolates KM10-BASS and CK2-BASS (Suppl. Fig. 2). Repeated attempts to culture the fungus were unsuccessful.

(b) *Spatiotemporal patterns of CFD and associated changes in seawater temperature*

Atoll-wide mean (\pm 1SE) CFD occurrence on Palmyra's forereef habitat equalled 0.1 ± 0.06 cases/m² of CCA in 2008 (total reef area surveyed in this habitat equalled 6,000 m²). Although cases of CFD were seen outside of our surveyed transects on the deeper (~10-15 m) reef terrace habitat, cases were rare in comparison; no CFD cases were documented within our surveyed transects on the shallow (<5 m) terrace or backreef habitats (total reef area surveyed in these two habitats equalled 5,800 m²). Within the forereef habitat, CFD occurrence displayed spatial heterogeneity ($Pseudo-F_{5,29} = 13.456$, $p < 0.0001$), with the central south forereef having the highest mean number of cases/m² of CCA (0.39 ± 0.10) in 2008 (Suppl. Table 1). Percent cover of CCA did not differ across forereef sites in 2008 ($Pseudo-F_{5,29} = 2.67$, $p = 0.06$, mean cover 23.5 %), and there was no relationship between CFD occurrence and CCA cover ($Pseudo-F_{1,29} = 0.311$, $p = 0.751$) (Suppl. Table 1).

Within the permanent backreef and reef terrace transects, mean CFD occurrence remained at 0 cases/m² of CCA in 2009 and 2010. However, within the permanent forereef transects, CFD mean occurrence increased approximately 14-fold from 0.10 cases/m² of CCA in 2008 to 1.37 cases/m² of CCA in late 2009 in association with the El Niño ocean warming event (Fig. 3, Suppl. Table 1). Seawater temperature had steadily increased over the latter part of 2009 and eventually peaked in November 2009, reaching 1.25°C and 1.51°C (satellite and forereef temperature observations, respectively) above the maximum long-term monthly climatological SST for Palmyra (Fig. 3). CFD occurrence increased in all permanent forereef transects during the El Niño event, with the central south forereef maintaining the highest levels (3.74 cases/m² of CCA, Suppl. Table 1). By March 2010, with a decrease in seawater temperatures, mean forereef CFD occurrence had returned to pre-El Niño levels (Fig. 3).

(c) *CFD vital rates during the El Niño event*

CFD lesions typically progressed in a radial manner across the surface of the CCA thallus, often crossing between individual CCA thalli but never spreading onto hard coral tissue (Suppl. Fig. 3). At time point zero, mean lesion surface area of the 13 CFD cases *in situ* was 108 mm² (± 25) (Suppl. Table 4). After one week, mean lesion surface area was 136 mm² (± 23), with a mean surface area progression rate of 3.5 mm²/day (± 2.3) and a mean linear progression rate of 2.4 mm/day (± 0.5). Across the entire four-week time period, the maximum surface area progression rate and linear progression rate of any single CFD lesion was 12.9 mm²/day and 6.5 mm/day, respectively (Suppl. Table 4).

(d) *Experimental effects of warming and acidification on CCA calcification and disease progression rates*

Exposure to elevated temperature and atmospheric *p*CO₂, designed to simulate ocean acidification (OA), reduced CCA net calcification rates, but this effect was dependent on fungal infection (Table 1). All CCA samples gained CaCO₃ mass in the ambient air treatments, while all samples lost mass in the elevated *p*CO₂ (lower pH) treatments (Fig. 4A). However, when exposed to both elevated *p*CO₂ and temperature, diseased CCA lost nearly twice as much mass as when exposed to simulated OA alone (significant interaction term, Table 1). Mass loss was not intensified for healthy CCA (Fig. 4A); net calcification rates in healthy CCA were significantly depressed only by elevated *p*CO₂ and not by elevated temperature (Table 1). Accordingly, calcification rates for diseased and healthy samples were statistically similar in all treatments, except for the simultaneous acidified and warmed conditions, in which diseased CCA lost 40% more mass than healthy CCA (t-test, df = 15, *p* = 0.0343). Visible lateral progression of the CFD lesion occurred only in the elevated temperature treatment in ambient CO₂ conditions where lesion size and lethality increased by 60% over one week (Fig. 4B).

4. DISCUSSION

Using a previously described crustose coralline algae fungal disease (CFD) [20], we demonstrate that ocean warming and acidification can have complex interactive effects on marine disease dynamics. These relationships are to be expected, as they reflect intricate relationships among the putative pathogen, host, and environment [39].

(a) *Identification of CFD-associated fungus*

Fungal pathogens are prevalent throughout the marine environment [40, 41], are commonly associated with the coral holobiont [42], and are known to infect tropical sea-fans [43, 44] and marine algae [45]. An inability to culture fungal isolates and a reliance on morphology for identification, however, have caused fungal isolates to be misclassified and their distribution underestimated [46]. Using histopathology of CFD samples from Palmyra Atoll and genetic sequencing of the associated fungus, we confirm a fungal infection of the CCA. While species-level identification was not possible, our phylogenetic analysis strongly suggests that the CFD fungus belongs to the subphylum *Ustilaginomycetes*, which consists of a large number of plant parasites, including strains of smut fungi [47]. Our methods, which allowed isolation of the fungus without an axenic culture, could be used to compare the fungus present in the Palmyra CFD lesions with fungi associated with suspected CFD lesions found on other reefs. These genetic approaches allow us to better interpret spatiotemporal dynamics of this disease on coral reefs and postulate their underlying mechanisms.

(b) *Disease dynamics and sea surface temperature*

Many fungal pathogens in animals and plants respond positively to elevated temperatures [48-51], and ocean warming is predicted to favour pathogens for many marine diseases [2]. At our

study location, CFD displayed a dramatic (14-fold) increase in occurrence on the forereef during an El Niño in association with sustained seawater temperatures well above the long-term climatological mean. Furthermore, we found experimentally that elevated temperature increases lateral expansion rates of CFD lesions. The positive relationship between temperature and CFD occurrence was likely the result of elevated temperatures increasing the virulence of the pathogen, and hence speeding disease progression, and causing physiological stress to the CCA host that ultimately reduced resistance to infection [52]. The prevalence of many coral diseases, such as white syndromes [5], atramentous necrosis [53], and black band disease [54] are also positively related to temperature. Interestingly, Vargas-Ángel (2010) [55] documented higher overall CCA disease occurrence at islands experiencing higher mean annual sea-surface temperature in a Pacific-wide survey of U.S.-affiliated coral reefs, further highlighting the importance of temperature in governing CCA disease dynamics.

While temperature variation provides a strong explanation for temporal variation in overall CFD occurrence at our study location, it does not adequately explain the spatial variation we documented at any one point in time. If higher temperatures cause both CFD occurrence and rates of lesion progression to increase, why was the disease almost exclusively limited to the forereef, where temperatures are lower than the shallow reef terrace [21]? There are several possible explanations. Host density is often a crucial factor determining the spatiotemporal distribution patterns of disease [39], with the prevalence of many coral and CCA diseases positively related to host abundance [5, 56, 57]. For example, Vargas-Ángel (2010) [55] found a positive relationship between island mean CCA cover and overall CCA disease occurrence at an archipelago scale across coral reefs of the U.S. Pacific, including the Pacific Remote Island Areas, the geopolitical region within which Palmyra resides. However, within Palmyra's forereef

habitat, CFD occurrence appeared to be independent of host abundance. Moreover, CCA cover peaks on Palmyra's shallow (<5 m) western reef terrace [10], where CFD was virtually absent; previous surveys at Palmyra have documented CFD on the terrace habitat, but again at deeper (~15 m) depths (Vargas-Ángel, NOAA, *pers. comm.*) where host cover is lower [10]. These findings suggest that host abundance alone does not explain the observed spatial variation in CFD occurrence at Palmyra. However, CCA species assemblages are known to vary spatially on reefs [58], and the peak of CFD occurrence on the forereef may simply reflect an increase in abundance of a preferred host species. While *in situ* species-specific assessments of CCA abundance would potentially resolve these issues, CCA taxonomy is difficult and requires microscopic examination, making it impossible in the field.

While potentially explaining between-habitat differences in disease occurrence, variation in host species abundance alone does not adequately explain the dramatic peak in CFD abundance on Palmyra's central south forereef. This CFD hotspot at Palmyra appears to be temporally stable, corroborating previous surveys conducted in 2006 [55]. The existence of this CFD hotspot at one site suggests that disease occurrence may be governed by external inputs of the pathogen, rather than by within-population transmission [44], or by an unmeasured extrinsic forcing. Palmyra's central south forereef is exposed to a particularly high level of lagoonal outflow during the change in tidal state (GJW *pers. obs.*), perhaps acting as a pathogen source and/or supplying more nutrient-rich waters that may enhance CFD establishment and progression, as has been shown for other fungal diseases on coral reefs [4]. Additionally, the south forereef of Palmyra has, on average, measurably higher seawater pH with less frequent or severe excursions than the north forereef or reef terrace [59]; our results indicate that the less acidic but warmer conditions characteristic of the southern forereef are most favourable for CFD occurrence. Regardless of the

mechanisms behind the fine-scale variations in CFD occurrence, it is clear that the disease is more abundant and virulent under elevated temperatures, suggesting that predicted increases in the frequency of temperature anomalies on coral reefs may result in more frequent CFD outbreaks.

(c) *Eco-physiological response of host to disease and climate change*

What will be the ecological consequences of increased CFD outbreaks on coral reefs? While elevated temperature increased overall CFD occurrence *in situ* and lateral rates of lesion expansion under experimental conditions, under the same experimental conditions elevated $p\text{CO}_2$ mediated these effects of temperature and slowed lesion expansion rates. While this suggests that future increases in the frequency of temperature anomalies will result in more frequent CFD outbreaks on coral reefs, the lowering of pH as a result of OA may actually slow overall spread of the disease across the reef landscape during such outbreaks. Importantly, however, while the lateral spread of CFD was not affected by reduced seawater pH and carbonate saturation state, all CCA thalli lost mass under OA conditions, suggesting net dissolution was occurring. For diseased thalli, these effects were exacerbated by warming.

Synergistic effects of ocean warming and acidification that together cause greater reduction in calcification of CCA than either stressor alone have been reported elsewhere [17-19, 60], but synergistic global climate change effects were only observed in this study when the CCA were also infected with the CFD fungus. Microboring organisms, or euendoliths, like fungi or cyanobacteria, burrow and erode carbonate at rates that can exceed biogenic CaCO_3 precipitation, leading to net dissolution of reef-building organisms [61, 62]. OA is expected to reduce resistance to euendolith penetration in both hermatypic corals and CCA by weakening

structural integrity of the CaCO_3 crystals [63], reducing skeletal density [64], and facilitating chemical dissolution [65, 66]. Further, colony formation is stimulated by natural reductions in pH, so OA has the potential to radically elevate abundance of marine fungi [67]. Not only can acidification weaken host resistance to bioerosion, but reduced saturation states and enhanced disease infestation of the CCA thallus could further accelerate corrosion. Thus synergistic interaction of pathogen infection, warming, and OA may exacerbate reef degradation under projected global climate change scenarios.

(d) *Conclusions*

Our study represents the first attempt to understand the interactive effects of two major global stressors, ocean warming and acidification, on disease dynamics on coral reefs. Using a fungal disease affecting crustose coralline algae (CFD), we show that while outbreaks of CFD should become more common on coral reefs as temperature anomalies become more frequent, ocean acidification may ameliorate lesion progression rates but still decrease overall survivorship of diseased hosts. The ecological consequences of such interactions are difficult to predict; however, it is clear that CFD possesses a tremendous capacity for lateral spread across the reef landscape during ocean warming events. Our results highlight the intricate nature of disease-host-environment interactions and the importance of adopting a multi-factor approach to modelling disease dynamics on coral reefs in order to accurately predict dynamics in a changing climate.

Acknowledgments We thank the U.S. Fish and Wildlife Service (USFWS) and The Nature Conservancy for granting access to Palmyra Atoll and providing logistical support. Funding was

provided by the National Geographic Society, the Gordon and Betty Moore Foundation, and a Victoria University of Wellington (VUW) Strategic Research Scholarship. We thank Rachel Morrison for edits to the manuscript and Brian Zgliczynski for logistical support. The majority of this research was conducted under the USFWS special use permits 12533-10010, 12533-11025, and 12533-12012. Scripps Institution of Oceanography is a member of the Palmyra Atoll Research Consortium (PARC). This is PARC publication number PARC-0098.

References

1. Ward J.R., Lafferty K.D. 2004 The elusive baseline of marine disease: Are diseases in ocean ecosystems increasing? *PLoS Biology* **2**(4), 542-547. (doi:e12010.1371/journal.pbio.0020120).
2. Harvell C.D., Mitchell C.E., Ward J.R., Altizer S., Dobson A.P., Ostfeld R.S., Samuel M.D. 2002 Ecology: Climate warming and disease risks for terrestrial and marine biota. *Science* **296**(5576), 2158-2162.
3. Aronson R.B., Precht W.F. 2001 White-band disease and the changing face of Caribbean coral reefs. *Hydrobiologia* **460**, 25-38.
4. Bruno J.F., Petes L.E., Harvell C.D., Hettinger A. 2003 Nutrient enrichment can increase the severity of coral diseases. *Ecology Letters* **6**(12), 1056-1061. (doi:10.1046/j.1461-0248.2003.00544.x).
5. Bruno J.F., Selig E.R., Casey K.S., Page C.A., Willis B.L., Harvell C.D., Sweatman H., Melendy A.M. 2007 Thermal stress and coral cover as drivers of coral disease outbreaks. *PLoS Biology* **5**(6), 1220-1227. (doi:e124 10.1371/journal.pbio.0050124).

- 423 6. Harvell D., Jordan-Dahlgren E., Merkel S., Rosenberg E., Raymundo L., Smith G., Weil
424 E., Willis B., Global Envriomn Facility C. 2007 Coral disease, environmental drivers, and the
425 balance between coral microbial associates. *Oceanography* **20**(1), 172-195.
- 426 7. Littler M.M., Littler D.S. 1984 Models of tropical reef biogenesis. *Phycological Research*
427 **3**, 324-364.
- 428 8. Price N. 2010 Habitat selection, facilitation, and biotic settlement cues affect distribution
429 and performance of coral recruits in French Polynesia. *Oecologia* **163**, 747-758.
- 430 9. Nyström M., Graham N.A.J., Lokrantz J., Norström A.V. 2008 Capturing the
431 cornerstones of coral reef resilience: linking theory to practice. *Coral Reefs* **27**(4), 795-809.
432 (doi:10.1007/s00338-008-0426-z).
- 433 10. Williams G.J., Smith J.E., Conklin E.J., Gove J.M., Sala E., Sandin S.A. 2013 Benthic
434 communities at two remote Pacific coral reefs: effects of reef habitat, depth, and wave energy
435 gradients on spatial patterns. *PeerJ* **1**, e81. (doi:DOI 10.7717/peerj.81).
- 436 11. Chisholm J.R. 2003 Primary productivity of reef-building crustose coralline algae.
437 *Limnology and Oceanography* **48**, 1376-1387.
- 438 12. Vargas-Angel B. 2010 Crustose coralline algal diseases in the U.S.-Affiliated Pacific
439 Islands. *Coral Reefs* **29**, 943-956.
- 440 13. Littler M.M., Littler D.S. 1995 Impact of CLOD pathogen on Pacific coral reefs. *Science*
441 **267**, 1356-1360.
- 442 14. Littler M.M., Littler D.S. 1997 Disease-induced mass mortality of crustose coralline
443 algae on coral reefs provide a rationale for conservation of herbivorous fish stocks. In *Proc 8th*
444 *Int Coral Reef Symp* (pp. 719-724).

- 445 15. Goreau T.J., Cervino J., Goreau M., Hayes R., Hayes M., Richardson L., Smith G.,
446 DeMeyer K., Nagelkerken I., Garzon-Ferrera J., et al. 1998 Rapid spread of diseases in
447 Caribbean coral reefs. *Revista De Biologia Tropical* **46**, 157-171.
- 448 16. Bulleri F. 2006 Duration of overgrowth affects survival of encrusting coralline algae.
449 *Marine Ecology Progress Series* **321**, 79-85.
- 450 17. Anthony K.R.N., Kline D.I., Diaz-Pulido G., Dove S., Hoegh-Guldberg O. 2008 Ocean
451 acidification causes bleaching and productivity loss in coral reef builders. *Proceedings of the*
452 *National Academy of Sciences of the United States of America* **105**(45), 17442-17446.
453 (doi:10.1073/pnas.0804478105).
- 454 18. Martin S., Gattuso J.P. 2009 Response of Mediterranean coralline algae to ocean
455 acidification and elevated temperature. *Global Change Biology* **15**(8), 2089-2100.
456 (doi:10.1111/j.1365-2486.2009.01874.x).
- 457 19. Johnson M.D., Carpenter R.C. 2012 Ocean acidification and warming decrease
458 calcification in the crustose coralline alga *Hydrolithon onkodes* and increase susceptibility to
459 grazing. *Journal of Experimental Marine Biology and Ecology* **434**, 94-101.
460 (doi:10.1016/j.jembe.2012.08.005).
- 461 20. Littler M.M., Littler D.S. 1998 An undescribed fungal pathogen of reef-forming crustose
462 coralline algae discovered in American Samoa. *Coral Reefs* **17**, 144.
- 463 21. Williams G.J., Knapp I.S., Maragos J.E., Davy S.K. 2010 Modeling patterns of coral
464 bleaching at a remote Central Pacific atoll. *Marine Pollution Bulletin* **60**, 1467-1476.
- 465 22. Williams G.J., Knapp I.S., Work T.M., Conklin E.J. 2011 Outbreak of *Acropora* white
466 syndrome following a mild bleaching event at Palmyra Atoll, Northern Line Islands, Central
467 Pacific. *Coral Reefs* **30**, 621.

- 468 23. Vargas-Angel B., Looney E.E., Vetter O.J., Coccagna E.F. 2011 Severe, widespread El
469 Nino-associated coral bleaching in the US Phoenix Islands. *Bulletin of Marine Science* **87**, 623-
470 638.
- 471 24. Kjer J., Debbab A., Aly A.H., Proksch P. 2010 Methods for isolation of marine-derived
472 endophytic fungi and their bioactive secondary products. *Nature protocols* **5**, 479 - 490.
- 473 25. Campbell E.L., Summers M.L., Christman H., Martin M.E., Meeks J.C. 2007 Global
474 gene expression patterns of *Nostoc punctiforme* in steady-state dinitrogen-grown heterocyst-
475 containing cultures and at single time points during the differentiation of akinetes and
476 hormogonia. *Journal of Bacteriology* **189**, 5247–5256.
- 477 26. Borneman J., Hartin R.J. 2000 PCR primers that amplify fungal rRNA genes from
478 environmental samples. *Applied and Environmental Microbiology* **66**, 4356.
- 479 27. Tamura K., Peterson D., Peterson N., Stecher G., Nei M., Kumar S. 2011 MEGA5:
480 Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and
481 maximum parsimony methods. *Molecular Biology and Evolution* **28**, 2731–2739.
- 482 28. Bass D., Howe A., Brown N., Barton H., Demidova M., Michelle H., Li L., Sanders H.,
483 Watkinson S.C., Willcock S., et al. 2007 Yeast forms dominate fungal diversity in the deep
484 oceans. *Proceedings of the Royal Society of London, Series B: Biological Sciences* **274**, 3069-
485 3077.
- 486 29. Tamura K., Nei M., Kumar S. 2004 Prospects for inferring very large phylogenies by
487 using the neighbor-joining method. *Proceedings of the National Academy of Sciences of the*
488 *United States of America* **101**, 11030–11035.

- 489 30. Gove J.M., Williams G.J., McManus M., Heron S., Sandin S.A., Vetter O.J., Foley D.
490 2013 Quantifying climatological ranges and anomalies for Pacific coral reef ecosystems. *PLoS*
491 *One* **8**(4), e61974.
- 492 31. Price N.N., Hamilton S.L., Tootell J.S., Smith J.E. 2011 Species-specific consequences of
493 ocean acidification for the calcareous tropical green algae *Halimeda*. *Marine Ecology-Progress*
494 *Series* **440**, 67-78. (doi:10.3354/meps09309).
- 495 32. IPCC. 2007 IPCC Fourth Assessment Report: Climate Change 2007. In *Climate Change* (
- 496 33. Valiela I. 1984 *Marine ecological processes*. New York, Springer-Verlag.
- 497 34. Davies P.S. 1989 Short term growth measurements of corals using an accurate buoyant
498 weighing technique. *Marine Biology* **101**(3), 389-395.
- 499 35. Lavigne H., Gattuso J.P. 2011 seacarb: seawater carbonate chemistry with R. (R package
500 version 2.4 ed, <http://CRAN.R-project.org/package=seacarb>).
- 501 36. Lueker T.J., Dickson A.G., Keeling C.D. 2000 Ocean pCO₂ calculated from dissolved
502 inorganic carbon, alkalinity, and equations for K-1 and K-2: validation based on laboratory
503 measurements of CO₂ in gas and seawater at equilibrium. *Mar Chem* **70**(1-3), 105-119.
504 (doi:10.1016/s0304-4203(00)00022-0).
- 505 37. Anderson M.J., Gorley R.N., Clarke K.R. 2008 PERMANOVA+ for PRIMER: Guide to
506 software and statistical methods. PRIMER-E: Plymouth, UK.
- 507 38. Anderson M.J. 2003 DISTLM forward: a FORTRAN computer program to calculate a
508 distance-based multivariate analysis for a linear model using forward selection. Department of
509 Statistics, University of Auckland, New Zealand.

- 510 39. Ostfeld R.S., Keesing F., Eviner V.T. 2008 *Infectious disease ecology: The effects of*
511 *ecosystems on disease and of disease on ecosystems*, Princeton University Press, New Jersey,
512 USA.; 506 p.
- 513 40. Rheinheimer G. 1992 *Aquatic Microbiology*. New York, Wiley.
- 514 41. Ramaiah N. 2006 A review on fungal diseases of algae, marine fishes, shrimps and
515 corals. *Indian Journal of Marine Sciences* **35**, 380-387.
- 516 42. Williams G.J., Work T.M., Aeby G.S., Knapp I.S., Davy S.K. 2011 Gross and
517 microscopic morphology of lesions in Cnidaria from Palmyra Atoll, Central Pacific. *Journal of*
518 *Invertebrate Pathology* **106**, 165-173.
- 519 43. Smith G.W., Ives L.D., Nagelkerken I.A., Ritchie K.B. 1996 Caribbean sea-fan
520 mortalities. *Nature* **383**(6600), 487-487.
- 521 44. Kim K., Harvell C.D. 2004 The rise and fall of a six-year coral-fungal epizootic. *The*
522 *American Naturalist* **164**(S5), S52-S63. (doi:10.1086/424609).
- 523 45. Raghukumar C. 1996 Zoosporic fungal parasites of marine biota. In *Advances in*
524 *zoosporic fungi* (ed. Dayal R.), pp. 61-83. New Delhi, M D Publications Pvt. Ltd.
- 525 46. Hawksworth D.L., Rossman A.Y. 1997 Where are all the undescribed fungi?
526 *Phytopathology* **87**(9), 888-891. (doi:10.1094/phyto.1997.87.9.888).
- 527 47. Bauer R., Begerow D., Oberwinkler E., Piepenbring M., Berbee M.L. 2001
528 Ustilaginomycetes. In *Systematics and Evolution* (eds. McLaughlin D., McLaughlin E., Lemke
529 P.), pp. 57-83, Springer Berlin Heidelberg.
- 530 48. Bemis D.A., Krahwinkel D.J., Bowman L.A., Mondon P., Kwon-Chung K.J. 2000
531 Temperature-sensitive strain of *Cryptococcus neoformans*: Producing hyphal elements in a feline
532 nasal granuloma. *Journal of Clinical Microbiology* **38**, 926-928.

49. Spitzer E.D., Keath E.J., Travis S.J., Painter A.A., Kobayashi G.S., Medoff G. 1990 Temperature-sensitive variants of *Histoplasma capsulatum* isolated from patients with acquired immunodeficiency syndrome. *Journal of Infectious Diseases* **162**, 258-261.
50. Hill R.A., Blankenship P.D., Cole R.J., Sanders T.H. 1983 Effects of soil moisture and temperature on preharvest invasion of peanuts by the *Aspergillus flavus* group and subsequent aflatoxin development. *Applied and Environmental Microbiology* **45**, 628-633.
51. Marin S., Sanchis V., Magan N. 1995 Water activity, temperature, and pH effects on growth of *Fusarium moniliforme* and *Fusarium proliferatum* isolates from maize. *Canadian Journal of Microbiology* **41**, 1063-1070.
52. Blanford S., Thomas M.B., Pugh C., Pell J.K. 2003 Temperature checks the Red Queen? Resistance and virulence in a fluctuating environment. *Ecology Letters* **6**(1), 2-5.
53. Jones R.J., Bowyer J., Hoegh-Guldberg O., Blackall L.L. 2004 Dynamics of a temperature-related coral disease outbreak. *Marine Ecology-Progress Series* **281**, 63-77.
54. Sato Y., Bourne D.G., Willis B.L. 2009 Dynamics of seasonal outbreaks of black band disease in an assemblage of *Montipora* species at Pelorus Island (Great Barrier Reef, Australia). *Proceedings of the Royal Society B-Biological Sciences* **276**(1668), 2795-2803. (doi:10.1098/rspb.2009.0481).
55. Vargas-Angel B. 2010 Crustose coralline algal diseases in the US-Affiliated Pacific Islands. *Coral Reefs* **29**(4), 943-956. (doi:10.1007/s00338-010-0646-x).
56. Aeby G.S., Williams G.J., Franklin E., Haapkyla J., Harvell C.D., Neale S., Page C.A., Raymundo L., Vargas-Angel B., Willis B., et al. 2011 Growth anomalies on the coral genera *Acropora* and *Porites* are strongly associated with host density and human population size across the Indo-Pacific. *PLoS One* **6**(2), e16887.

57. Myers R.L., Raymundo L.J. 2009 Coral disease in Micronesian reefs: a link between disease prevalence and host abundance. *Diseases of Aquatic Organisms* **87**(1/2), 97-104. (doi:10.3354/dao02139).
58. Dethier M.N., Paull K.M., Woodbury M.M. 1991 Distribution and thickness patterns in subtidal encrusting algae from Washington. *Botanica Marina* **34**, 201-210.
59. Price N.N., Martz T.R., Brainard R.E., Smith J.E. 2012 Diel variability in seawater pH relates to calcification and benthic community structure on coral reefs. *Plos One* **7**(8), e43843. (doi:10.1371/journal.pone.0043843).
60. Diaz-Pulido G., Anthony K.R.N., Kline D.I., Dove S., Hoegh-Guldberg O. 2012 Interactions between ocean acidification and warming on the mortality and dissolution of coralline algae. *Journal of Phycology* **48**(1), 32-39. (doi:10.1111/j.1529-8817.2011.01084.x).
61. Godinot C., Tribollet A., Grover R., Ferrier-Pages C. 2012 Bioerosion by euendoliths decreases in phosphate-enriched skeletons of living corals. *Biogeosciences* **9**(7), 2377-2384. (doi:10.5194/bg-9-2377-2012).
62. Tribollet A., Payri C. 2001 Bioerosion of the coralline alga *Hydrolithon onkodes* by microborers in the coral reefs of Moorea, French Polynesia. *Oceanologica Acta* **24**(4), 329-342. (doi:10.1016/s0399-1784(01)01150-1).
63. Ragazzola F., Foster L.C., Form A., Anderson P.S.L., Hansteen T.H., Fietzke J. 2012 Ocean acidification weakens the structural integrity of coralline algae. *Global Change Biology*. (doi:10.1111/j.1365-2486.2012.02756.x).
64. Hernandez-Ballesteros L.M., Elizalde-Rendon E.M., Carballo J.L., Carricart-Ganivet J.P. 2013 Sponge bioerosion on reef-building corals: Dependent on the environment or on skeletal

density? *Journal of Experimental Marine Biology and Ecology* **441**, 23-27.

(doi:10.1016/j.jembe.2013.01.016).

65. Tribollet A., Atkinson M.J., Langdon C. 2006 Effects of elevated pCO₂ on epilithic and endolithic metabolism of reef carbonates. *Global Change Biology* **12**, 2200-2208.

(doi:10.1111/j.1365-2486.2006.01249.x).

66. Wisshak M., Schonberg C.H.L., Form A., Freiwald A. 2012 Ocean acidification accelerates reef bioerosion. *Plos One* **7**(9). (doi:e4512410.1371/journal.pone.0045124).

67. Krause E., Wichels A., Giménez L., Gerdt G. 2013 Marine fungi may benefit from ocean acidification. *Aquat Microb Ecol* **69**(1), 59-67.

597 Table 1. Two-way ANOVA results for calcification rates of diseased and healthy CCA crusts (n
 598 = 8 per treatment) immersed in the ocean acidification ('CO₂ enrichment') and ambient
 599 conditions across duplicate flow-through seawater tables ('Table') nested within a warming El
 600 Niño ('Temperature') or a seasonal average scenario.

601

CCA state	Source	DF	F	p
<i>Healthy, no lesions</i>	CO ₂ enrichment	1	64.327	< 0.0001
	Temperature	1	0.018	0.896
	CO ₂ x Temp.	1	0.050	0.824
	Table [Temp.]	2	1.667	0.208
<i>Diseased, lesions present</i>	CO ₂ enrichment	1	163.153	< 0.0001
	Temperature	1	3.008	0.095
	CO ₂ x Temp.	1	4.593	0.042
	Table [Temp.]	2	0.396	0.677

602

603

604

605

606

607

608

609

610

Figure legends

Figure 1. Location of Palmyra Atoll and the permanent monitoring sites established in 2008 (1-6 = shallow terrace; 7-12 = forereef).

Figure 2. Field signs of coralline fungal disease (CFD) (A). Fungal infection appears as a blue-black lesion (arrows). Days-old exposed substrate becomes colonised by microalgae and turf algae (1) and appears bleached white when freshly exposed (2), while the CCA tissue remains pink and healthy on the leading edge of the lesion (3). Bar = 1 cm. (B) Appearance of isolated fungal hyphae associated with CFD (1000x magnification using light microscopy). Bar = 15 μ m. (C) Section of a coralline alga infected with CFD and positively confirmed as a fungal infection using Grocott's methenamine silver. Note the fungal hyphae invading the algal thallus and conceptacles (arrows). Cu = cuticle. Bar = 30 μ m.

Figure 3. Representative *in situ* temperatures at Palmyra Atoll at 10 m on the forereef and sea-surface temperatures (SST) from satellite-derived sources during 2008, 2009, and 2010 and the associated change in coralline fungal disease (CFD) occurrence (forereef-wide mean number of cases/m² of CCA are shown by black arrows).

Figure 4. (A) Mean (\pm SE) calcification rate for diseased and healthy CCA in experimental aquaria (n = 12). Change in weight shown as mg CaCO₃/week for each thallus. Asterisk indicates when response within treatment differs significantly between diseased and healthy specimens

632 (Table 2). (B) Mean (\pm SE) lesion lateral expansion rate for diseased CCA in experiments. Using
633 the ambient air x 28°C treatment as a control value, the asterisk indicates a significant effect of
634 elevated temperature (Dunn's, $Z = 3.156$, $p = 0.0048$).